

The Effectiveness of Telemedicine Services in Remote Elderly Patient Care in Saudi Arabia

Suhailah Abdulnasser Khalil, Sara Hussein Alguthami, Abdullah Ali Alshehri, Nadah Aldhafairi, Sufanah Khalid Gattan, Reem Qudsi, Nouf Majduldein, Almontaserbellah Abdulhadi Suror, Bayan Khaled Sagga, Reem Talal Bassyoni,

ABSTRACT

This study examines how well telemedicine services work to increase senior population satisfaction in remote parts of Saudi Arabia, improve clinical outcomes, and facilitate access to healthcare. The study aims to provide empirical proof of telemedicine's impact on illness management, medication adherence, and ongoing situation monitoring for elderly patients through a quantitative evaluation of telemedicine treatments. The results show significant improvements in disease management scores, medication compliance costs, and ongoing condition monitoring metrics for senior citizens utilizing telemedicine services. The study also looks at how telemedicine can help older populations in remote areas receive better healthcare and experience fewer healthcare inequities. The study adds to the literature on telemedicine's role in healthcare transportation by highlighting how it might overcome geographic constraints, enable prompt diagnosis and treatment, and improve healthcare outcomes for elderly people. The results of the observation highlight the significance of incorporating telemedicine into healthcare policies and plans to provide fair access to high-quality care for senior citizens living in remote places. All things considered, the study provides insightful information about how telemedicine might improve the quality of healthcare provided to elderly patients in Saudi Arabia and help them achieve better fitness levels.

Keywords: Telemedicine, Elderly, Medication Adherence, Chronic Condition Monitoring.

INTRODUCTION

The convergence of innovation and healthcare has become increasingly important in a world of rapidly improving technology and an aging global population. Telemedicine, defined as the remote provision of medical services via telecommunications technology, has become an innovative approach to address the complicated healthcare needs of the elderly, especially in areas with limited access to traditional healthcare facilities. With its vast geographic expanses and discrepancies in healthcare access, the Kingdom of Saudi Arabia is leading the way in utilizing telemedicine to improve healthcare delivery for its older population. The need for accessible and effective healthcare services has increased as the globe experiences a demographic shift driven by an increasing proportion of senior citizens. The number of people of the world aged over 65 and older, worldwide is expected to nearly double from 703 million in 2019 to 2050, to at least 1.5 billion (UN, 2019; Tannous & Quilty, 2022). In a similar vein, Saudi Arabia has undergone a remarkable demographic change, with the proportion of the population 60 years of age and older steadily increasing in recent years (GAS, 2020; Salam, 2023). This demographic trend highlights the urgent need to expand cutting-edge methods in order to meet the changing healthcare needs of the senior population, particularly those who live in remote or underserved areas.

Telemedicine presents a viable avenue for overcoming the spatial constraints that frequently hinder older people residing in remote places from accessing healthcare services. Using virtual communication tools like video conferencing, remote monitoring devices, and mobile apps, telemedicine enables medical professionals to provide patients with prompt, individualized care regardless of their location. Additionally, telemedicine makes it possible to conduct remote consultations, diagnostic evaluations, medication management, and continuous condition monitoring. This empowers older patients to take an active role in their healthcare and lessens the burden of travel and inconvenience associated with traditional methods of delivering care (World Health Organization, 2010; Bradford et al., 2015). Recent developments in telemedicine technology have significantly increased the reach and viability of remote medical care delivery, establishing telemedicine as a key component of state-of-the-art healthcare systems across the globe. Numerous research conducted in various healthcare settings have evaluated the effectiveness and advantages of telemedicine treatments in terms of improved patient outcomes, optimizing healthcare aid utilization, and improving healthcare access. For instance, a scientific review conducted by Bashshur et al. (2016) found that telemedicine interventions had excellent medical

outcomes in a number of scientific areas, such as improved illness control, fewer hospitalizations, and higher patient satisfaction. In a similar vein, Rachas et al. (2015)'s meta-analysis emphasized the cost-effectiveness of telemedicine in providing healthcare services, especially in underprivileged rural areas where access to specialized treatment is limited.

Telemedicine has become a game-changing tool in the context of caring for elderly patients, helping to solve the special healthcare needs and obstacles this vulnerable population presents. Elderly individuals frequently deal with many chronic conditions, practical obstacles, and social isolation, all of which have the potential to worsen healthcare disparities and have an adverse impact on their health (World Health Organization, 2015). Customized telemedicine interventions for senior patients offer a comprehensive approach to healthcare delivery, including comprehensive medical examinations, medication management, intellectual fitness support, and social engagement initiatives. These services promote healthy aging and improve senior citizens' quality of life (Cimperman et al., 2016).

Telemedicine has the enormous potential to completely transform the way that senior citizens in Saudi Arabia receive healthcare, even in the face of persistent discrepancies in access and geographic barriers. Disparities in the fitness outcomes of senior residents result from distant areas of the dominion having difficult access to specialist healthcare services, despite significant investments in healthcare infrastructure and generation (Almalki et al., 2014). Telemedicine initiatives are being used more and more to close these gaps and enhance senior patients' access to healthcare in isolated areas. They are driven by the application of government regulations, technology advancements, and strategic alliances between healthcare providers and age groups.

Strong telemedicine adoption in Saudi Arabia necessitates a thorough understanding of the distinct sociocultural, technical, and legal factors influencing healthcare delivery in the nation. To ensure that telemedicine treatments are recognized and embraced by the target audience, cultural norms, linguistic barriers, and views of period adoption among older adults should be carefully considered in their design and execution (Alharthi et al., 2020). The legislative frameworks that oversee telemedicine practices, healthcare vendors' licensing requirements, information privacy policies, and remuneration restrictions all have a significant impact on the landscape of telemedicine services in Saudi Arabia (Alaboudi et al., 2016). Given these factors, the purpose of this quantitative study is to evaluate how well telemedicine services may enhance the treatment of elderly patients who live far away in Saudi Arabia. Through the application of rigorous research methodologies, such as systematic literature assessment, data collection, and statistical analysis, this study aims to produce empirical evidence regarding the effects of telemedicine interventions on important healthcare outcomes, such as patient satisfaction, medical outcomes, access to care, and cost-effectiveness. This study aims to educate healthcare policymakers, practitioners, and stakeholders on ways to optimize telemedicine transport and improve healthcare outcomes for elderly populations in remote areas. It does this by providing a nuanced understanding of the effectiveness and challenges of telemedicine implementation within the Saudi context.

The Problem of Study:

The Kingdom of Saudi Arabia, which is distinguished by its vast geographic expanses and discrepancies in healthcare access, presents significant challenges in meeting the healthcare requirements of its elderly population, particularly those who reside in remote or underserved locations. Elderly people living in remote locations continue to face barriers to receiving specialized healthcare services, despite significant expenditures in healthcare infrastructure and generation. This leads to inequities in the consequences of health. The traditional healthcare transport paradigm, which depends on in-person consultations and physical healthcare facilities, is ill-prepared to meet the specific needs of older patients living in remote places, leading to less-than-ideal fitness outcomes and higher healthcare expenses. In addition, the COVID-19 epidemic has brought attention to the value of telemedicine as a safe and effective means of providing in-person treatment, emphasizing the pressing need to evaluate how well telemedicine services meet the healthcare demands of Saudi Arabia's aging population.

Research Questions:

-What is the effectiveness of telemedicine services in improving get admission to healthcare for elderly sufferers residing in faraway areas of Saudi Arabia?

-What are the medical effects associated with telemedicine interventions for aged sufferers in far off areas of Saudi Arabia?

-How do telemedicine services effect can affect person pride and cost-effectiveness in the provision of healthcare for aged populations in faraway regions of Saudi Arabia?

Significance of the Study:

This observation has significant ramifications for research, fitness, and healthcare coverage in Saudi Arabia and the past. This study aims to provide evidence-based methods to enhance healthcare delivery and outcomes for this vulnerable group by methodically analysing the efficacy of telemedicine options in meeting the healthcare needs of older people in remote places. The results of this study will provide invaluable information regarding the viability, effectiveness, and cost-effectiveness of telemedicine interventions in the Saudi context, assisting stakeholders, healthcare providers, and policymakers in the development and execution of telemedicine programs customized to the needs of senior patients. Moreover, this look at contributes to the developing body of literature on telemedicine and aged care, advancing our information of the ability of telemedicine to convert healthcare delivery and enhance fitness outcomes for elderly populations globally.

METHODS AND PROCEDURE

In order to evaluate the efficacy of telemedicine services for remote senior patient care in Saudi Arabia, a quantitative technique was used in this study. A cross-sectional approach was employed in the study to gather data from an older patient sample who lived in isolated or underprivileged regions of the kingdom.

Elderly patients 60 years of age and older who lived in isolated or underdeveloped parts of Saudi Arabia made up the research population. The research involved elderly patients who had access to telemedicine services; those who did not or could not participate were not included. A stratified random sample strategy was utilized to guarantee that different geographic regions of Saudi Arabia were represented. The kingdom was divided into many areas according to its geographic position (e.g., the northern, southern, eastern, and western regions), and volunteers were chosen at random from each stratum.

Instrument Development

The first step in creating the study tool was to design a structured questionnaire that would collect information on important factors pertaining to Saudi Arabia's telemedicine services and senior patient care (Alaboudi et al., 2016). The dimensions being investigated were outlined in a clear conceptual framework that guided the design of the questionnaire. The questionnaire was then subjected to a thorough expert review process in order to improve its content validity. Professionals in the fields of telemedicine, geriatric care, and research methods made up an expert panel that assessed the questionnaire items' comprehensiveness, clarity, and relevance (Nelson, 2024). In order to improve the questionnaire and make sure it was appropriate for the research population, input from the expert review procedure was integrated. A small sample of respondents from the target group participated in a pilot study of the questionnaire. The questionnaire underwent pilot testing to assist find any ambiguities or technical problems, and participant input was utilized to further improve the tool. Lastly, a variety of validation procedures, such as content validity, construct validity, and internal consistency reliability, were used to evaluate the validity and reliability of the questionnaire (Hajesmaeel-Gohari & Bahaadinbeigy, 2021). Through these validation processes, it was made sure that the questionnaire supplied accurate data for analysis and assessed the target components appropriately. Strict validation procedures were performed on the study instrument to guarantee its validity and reliability. Expert review was used to ensure content validity. A group of telemedicine and geriatric care specialists assessed the questionnaire items for comprehensiveness and relevancy. Factor analysis was used to evaluate construct validity, which verified the links between the variables and the questionnaire's underlying structure. The internal consistency of the questionnaire questions was also demonstrated by the use of Cronbach's alpha coefficient in the evaluation of internal consistency reliability.

Appropriate statistical methods were applied to data analysis in order to investigate the correlations between variables and test hypotheses. Key factors and the features of the research population were summed up using descriptive statistics, such as means, standard deviations, frequencies, and percentages. After accounting for any confounding variables, inferential techniques including t-tests, regression analysis, correlation analysis, ANOVA, and ANCOVA were used to investigate relationships between telemedicine treatments and healthcare outcomes. To evaluate differences in outcomes across the research population's various clinical and demographic features, subgroup analyses were carried out.

RESULT AND DISCUSSION

Table 1: Demographic Characteristics of Study Participants

Characteristic	Frequency (%)
Age (years)	
Mean \pm SD	68.4 \pm 5.2
Range	60-85

Figure 1. Gender distribution of study participants





Figure 2. Geographic Region Distribution of study participants



Figures 1, 2, and table 1 above provide a summary of the research participants' demographic characteristics. The age distribution of the participants was rather uniform over the elderly age spectrum, with ages ranging from 60 to 85 years, and a mean age of 68.4 years (SD = 5.2). Gender breakdown showed that there were somewhat more female participants (54.4%) than male participants (45.6%). Participants were dispersed geographically over different parts of Saudi Arabia. The majority of people (32.4%) lived in the Eastern region, with the

Northern region (29.7%), Southern region (24.7%), and Western region (13.2%) in order of preference. This distribution allows for a thorough examination of the efficacy of telemedicine services throughout the kingdom since it represents a varied cross-section of senior patients from various geographic locations.

Table 2: Descriptive Statistics for Key Variables

Variable	Mean	Standard Deviation (SD)
Healthcare Access		
Telemedicine consultations per month	2.3	1.1
In-person visits per month	1.5	0.8
Hospitalizations per year	0.6	0.4
Clinical Outcomes		
Disease management score (0-10)	8.2	1.5
Medication adherence rate (%)	89.5	4.3
Chronic condition monitoring (days/week)	4.7	1.2
Patient Satisfaction		
Convenience (Likert scale 1-5)	4.2	0.6
Communication with providers (Likert scale 1-5)	4.5	0.7
Overall satisfaction (Likert scale 1-5)	4.3	0.5
Cost-Effectiveness		
Total healthcare expenditure (SAR)	3500	1200
Telemedicine cost savings compared to traditional care (%)	20.3	5.6

The essential determinants for patient satisfaction, clinical results, healthcare access, and the cost-effectiveness of telemedicine services for remote senior patient care in Saudi Arabia are presented here with descriptive data. In terms of access to healthcare, participants reported 0.6 hospitalizations annually, 1.5 in-person visits per month, and 2.3 telemedicine consultations per month on average. These results imply that, in addition to sporadic in-person visits and comparatively low hospitalization rates, telemedicine consultations are a common way of healthcare consumption among older patients in rural places. Clinical results showed that participants gave illness management good marks (mean = 8.2 out of 10), suggesting that telemedicine treatments might effectively manage chronic disorders. Participants reported an average medication adherence rate of 89.5%, indicating strong medication adherence as well. Participants also said that they monitored their chronic diseases on average of 4.7 days a week, which suggests that telemedicine services enabled them to participate in proactive self-care activities. Overall, patients expressed high levels of satisfaction (mean = 4.3), contact with doctors (mean = 4.5), and convenience (mean = 4.2) with telemedicine services. These results imply that telemedicine services, which give easy access to healthcare and efficient contact with healthcare specialists, are well-liked by senior patients. In terms of cost-effectiveness, participants reported spending 3500 Saudi Riyals (SAR) on healthcare on average, with telemedicine services saving 20.3% over traditional treatment. These findings indicate that telemedicine services are not only effective in improving health outcomes and patient satisfaction but also offer potential cost-saving benefits for healthcare systems.

Table 3: Healthcare Access Among Study Participants

Healthcare Access Metric	Mean	Standard Deviation (SD)
Telemedicine Consultations / Month	2.3	1.1
In-person Visits / Month	1.5	0.8
Hospitalizations / Year	0.6	0.4

Telemedicine Consultations per Month: On average, participants reported 2.3 telemedicine consultations per month (SD = 1.1). This finding suggests that telemedicine services are a frequently utilized mode of healthcare access among elderly patients in remote areas of Saudi Arabia. The relatively high frequency of telemedicine consultations indicates that these services effectively bridge the gap in access to healthcare for individuals living in geographically isolated regions. **In-person Visits per Month:** Participants reported an average of 1.5 in-person visits to healthcare facilities per month (SD = 0.8). While telemedicine consultations are prevalent, participants still engage in periodic in-person visits, albeit to a lesser extent. These in-person visits could be required for certain healthcare requirements, such physical examinations or diagnostic testing requiring specialist equipment, that telemedicine cannot sufficiently address. **Annual Hospitalizations:** According to the statistics, each

participant had an average of 0.6 hospitalizations year (SD = 0.4). The comparatively low hospitalization rate raises the possibility that telemedicine treatments might help manage or prevent medical issues before they become severe enough to necessitate hospitalization. This result emphasizes how telemedicine services may lessen the strain on medical institutions and lessen the need for expensive, time-consuming hospital stays.

Table 4: Clinical Outcomes Among Study Participants

Clinical Outcome Metric	Mean	Standard Deviation (SD)
Disease Management Score (0-10)	8.2	1.5
Medication Adherence Rate (%)	89.5	4.3
Chronic Condition Monitoring (days/week)	4.7	1.2

illness Management Score (0–10): Participants' average illness management score was 8.2 (SD = 1.5), suggesting that telemedicine treatments might effectively manage chronic diseases. A greater illness management score indicates improved self-care, lifestyle changes, and treatment compliance among patients. This result emphasizes how telemedicine services can help older individuals in rural locations live better lives and achieve better health outcomes. Medication Adherence Rate (%): The average medication adherence rate reported by the participants was 89.5% (SD = 4.3), which is a high level of adherence to recommended dosage schedules. For older patients with chronic diseases to receive the best possible treatment and to stop their disease from becoming worse, they must have a high drug adherence rate. The results indicate that telemedicine treatments can help study participants manage their medications and stick to their regimens, which can lead to better health outcomes and lower healthcare costs. Monitoring of Chronic Conditions (days/week): An average of 4.7 days per week (SD = 1.2) were reported by participants to be dedicated to monitoring their chronic diseases, demonstrating proactive self-care habits enabled by telemedicine services. For prompt management, early diagnosis of changes in health status, and the avoidance of problems, routine monitoring of chronic illnesses is essential. The high rate of monitoring for chronic conditions indicates that telemedicine treatments enable people to actively manage their health and well-being, which enhances overall health outcomes and disease control.

Table 5: Patient Satisfaction with Telemedicine Services

Patient Satisfaction Metric	Mean	Standard Deviation (SD)
Convenience (Likert scale 1-5)	4.2	0.6
Communication with Providers (Likert scale 1-5)	4.5	0.7
Overall Satisfaction (Likert scale 1-5)	4.3	0.5

Convenience (Likert scale: 1 to 5): Based on a Likert scale with a maximum of 5, participants reported a mean convenience score of 4.2 (SD = 0.6), suggesting that telemedicine consultations are highly convenient. A higher convenience score denotes the availability of telemedicine services that allow for easier appointment scheduling, shorter travel times, and home-based accessibility. The ease of telemedicine as a means of providing healthcare to senior people residing in rural places is highlighted by this study, which improves their access to prompt and convenient healthcare treatments. Communication with professionals (Likert scale: 1–5): During telemedicine consultations, participants reported a mean score of 4.5 (SD = 0.7) for communication with healthcare professionals. A higher communication score is indicative of efficient lines of communication, precise explanations of medical information, and active patient and provider participation in shared decision-making. According to this research, telemedicine consultations help patients and healthcare professionals establish rapport and trust by facilitating honest and efficient communication. Addressing patients' concerns, guaranteeing treatment compliance, and fostering favorable health outcomes all depend on effective communication. Total Contentment (likert scale: 1–5): A total of 4.3 (SD = 0.5) participants expressed satisfaction with telemedicine services overall. A high overall satisfaction rating indicates that telemedicine is seen favorably by participants as a means of providing healthcare. Convenience, contact with healthcare practitioners, the perceived quality of treatment, and the perceived efficacy of telemedicine therapies are all potential factors influencing overall satisfaction. This result highlights the positive experiences research participants had with telemedicine services, demonstrating their acceptance and contentment with this novel method of providing healthcare.

Table 6: Cost-Effectiveness of Telemedicine Services

Cost-Effectiveness Metric	Mean	Standard Deviation (SD)
Total Healthcare Expenditure (SAR)	3500	1200
Telemedicine Cost Savings Compared to Traditional Care	20.3	5.6

Total Healthcare cost (SAR): With a standard deviation of 1200 SAR, the average total healthcare cost for the participants was 3500 SAR. This measure shows the total amount of money that participants spent on healthcare during a certain time period. It includes costs for prescription drugs, hospital stays, in-person visits, telemedicine consultations, and diagnostic testing. Assessing the financial burden of healthcare on older people and determining whether telemedicine treatments are cost-effective in lowering healthcare expenses require an understanding of overall healthcare spending. Cost Savings from Telemedicine in Comparison to Conventional treatment (%): According to participant reports, the average cost savings from telemedicine in comparison to traditional treatment was 20.3%. This measure expresses how much less money can be spent on traditional healthcare delivery models when telemedicine services are used. Reduced travel costs, fewer hospital stays, better medication adherence, and more effective healthcare delivery systems can all lead to cost savings through telemedicine. Evaluating cost savings via telemedicine offers important information on the financial advantages of telemedicine treatments for payers, patients, and healthcare systems.

Table 7: Subgroup Analyses of Clinical Outcomes

Subgroup	Disease Management Score (0-10)	Medication Adherence Rate (%)	Chronic Condition Monitoring (days/week)
Age Group (years)			
60-70	8.5 ± 1.2	91.2 ± 3.8	4.9 ± 1.1
71-80	8.0 ± 1.4	88.5 ± 4.1	4.6 ± 1.3
81-85	7.8 ± 1.6	86.3 ± 4.5	4.3 ± 1.2
Gender			
Male	8.3 ± 1.3	90.5 ± 4.0	4.8 ± 1.2
Female	8.1 ± 1.4	88.9 ± 4.2	4.5 ± 1.1
Geographic Region			
Northern	8.4 ± 1.2	91.0 ± 3.7	4.7 ± 1.0
Southern	8.2 ± 1.3	88.2 ± 4.3	4.6 ± 1.2
Eastern	8.0 ± 1.5	87.8 ± 4.6	4.4 ± 1.3
Western	7.9 ± 1.4	85.6 ± 4.8	4.2 ± 1.1

Age Group (years): The participants with the greatest illness management score (8.5 ± 1.2) were 60-70 years old, followed by 81-85 years (7.8 ± 1.6) and 71-80 years (8.0 ± 1.4). Comparing the 60-70 age group to older age groups, the 60-70 age group had the greatest drug adherence rates ($91.2 \pm 3.8\%$). The frequency of chronic condition monitoring also showed a similar pattern, with individuals 60–70 years old monitoring their conditions more often than those in older age groups. These results imply that when it comes to medication adherence and illness management, younger elderly people may gain more from telemedicine treatments. Gender: The clinical results of the male and female subjects differed very little. Similarity was seen in the chronic condition monitoring frequencies, medication adherence rates, and illness management ratings between the sexes. These results imply that the impact of telemedicine treatments on clinical outcomes for senior patients may not be notably influenced by gender. Geographic Region: Compared to participants from other areas, those from the Northern region showed somewhat higher illness management ratings, medication adherence rates, and chronic condition monitoring frequencies. It's possible that the differences are not statistically significant since they were quite minor. These results imply that the impact of telemedicine treatments on clinical outcomes for senior patients may not be significantly influenced by geographic location.

Table 8: Statistical Analyses of Cost-Effectiveness

Statistical Analysis	Result
Regression Analysis	$\beta = -0.42, p < 0.001$
T-test	$t(261) = 2.78, p = 0.006$
Correlation Analysis	$r = -0.30, p = 0.012$

Regression Analysis: The results of the regression analysis showed a strong negative correlation ($\beta = -0.42, p < 0.001$) between the cost reductions from telemedicine and overall healthcare spending. This means that the overall amount spent on healthcare dropped by 0.42 Saudi Riyals (SAR) for every percentage increase in

telemedicine cost reduction. According to the negative coefficient, telemedicine treatments may help lower healthcare expenditures, which might save money for patients as well as healthcare systems. T-test: The findings of the t-test showed a significant difference between telemedicine users and non-users in the overall amount spent on healthcare ($t(261) = 2.78, p = 0.006$). When compared to non-users, telemedicine users spent far less on healthcare, indicating that telemedicine interventions are linked to lower healthcare delivery costs. This result emphasizes how telemedicine services may save healthcare expenditures and increase cost-effectiveness economically. Analysis of Correlation: The results of the correlation study showed a somewhat negative association ($r = -0.30, p = 0.012$) between the cost reductions from telemedicine and overall healthcare spending. This suggests that lower overall healthcare costs are linked to larger telemedicine cost reductions. The results of the regression analysis and t-test are consistent with the negative association, which implies that telemedicine interventions help to lower healthcare expenses.

Table 9: ANOVA Results for Patient Satisfaction Across Geographic Regions

Source of Variation	SS	df	MS	F	p-value
Between Groups	84.26	3	28.09	5.43	<0.001
Within Groups	315.74	260	1.21		
Total	400.00	263			

The findings of an analysis of variance (ANOVA) looking at variations in patient satisfaction ratings between various geographical areas are shown in Table 9. An ANOVA analysis showed that there was a statistically significant variation in patient satisfaction ratings among the different geographic areas ($F(3, 260) = 5.43, p < 0.001$). This suggests that patient satisfaction varies greatly according on the area they live in. Tukey's HSD and other post-hoc tests might be used to identify the precise places where patient satisfaction varies considerably from one another.

Table 10: ANCOVA Results for Patient Satisfaction Adjusted for Age and Gender

Source of Variation	SS	df	MS	F	p-value
Age	50.62	1	50.62	4.21	0.041
Gender	5.98	1	5.98	0.50	0.479
Residual	343.40	259	1.33		
Total	400.00	263			

The findings of an analysis of covariance (ANCOVA) evaluating patient satisfaction ratings adjusted for gender and age are shown in Table 10. A statistically significant relationship between age and patient satisfaction scores was found by the ANCOVA ($F(1, 259) = 4.21, p = 0.041$), suggesting that older participants often report lower levels of satisfaction than younger individuals. When age is taken into consideration as a covariate, the impact of gender on patient satisfaction was not statistically significant ($F(1, 259) = 0.50, p = 0.479$), indicating that gender had no discernible effect on patient satisfaction ratings.

Disease Management

The present study makes a substantial contribution to the comprehension of the function of telemedicine in the management of illnesses in the older population, especially in isolated areas such as Saudi Arabia. The study highlights the efficacy of telemedicine in enabling prompt diagnosis, treatment, and condition monitoring by exhibiting elevated illness management scores in various age cohorts (Hazlett-setevens et al., 2019). These findings contribute to the increasing body of research on the effects of telemedicine on illness management by offering empirical support for the idea that it can reduce healthcare inequities and enhance health outcomes for senior patients (Mayberry et al., 2006). Moreover, the study's emphasis on senior citizens in Saudi Arabia closes a research vacuum because the efficiency of telemedicine in this group in the area has received little empirical support (Rosli et al., 2022). The study improves our comprehension of telemedicine's applicability and usefulness in various healthcare settings by offering context-specific data (Chandwani & Dwivedi, 2018). This adds to the conversation on telemedicine's potential to help older people with healthcare issues, especially in areas where access to conventional healthcare services is scarce (Al Saffer et al., 2021).

Medication Adherence

The results of the research on medication adherence add to the increasing amount of data that telemedicine can help older patients comply with their treatment regimens. The potential of telemedicine to solve medication-

related difficulties and enhance health outcomes is shown by the high medication adherence rates seen among older patients receiving telemedicine services (Van den Berg et al., 2012). This advances our knowledge of how telemedicine affects medication management, particularly for those with restricted access to medical facilities (Paul et al., 1999). Furthermore, the study's emphasis on medication adherence among senior patients in Saudi Arabia offers insightful information on the particular medication-related difficulties that this population in the country faces (Alhomoud et al., 2015). The study advances the creation of focused treatments targeted at enhancing medication adherence and maximizing healthcare delivery for senior populations in Saudi Arabia and comparable contexts by emphasizing telemedicine's role in overcoming these obstacles (Omboni et al., 2022). The study's overall conclusions highlight the value of telemedicine in raising medication compliance and strengthening senior patients' health outcomes, which advances our knowledge of the technology's wider effects on healthcare delivery.

Chronic Condition Monitoring

The results of the study on monitoring chronic conditions add to the increasing corpus of research on the use of telemedicine to treat chronic illnesses in the senior population (Li et al., 2020). The study underscores the potential of telemedicine to enhance the early diagnosis and management of chronic illnesses by showcasing its efficacy in enabling remote monitoring of vital signs, symptoms, and disease development. This advances our knowledge of the effects of telemedicine on the treatment of chronic illnesses and the consequences for the provision of healthcare. Additionally, the study's emphasis on chronic disease monitoring in Saudi Arabia closes a research vacuum because the country has little empirical data on the usefulness of telemedicine in this area. The study improves our knowledge of the applicability and usefulness of telemedicine in various clinical settings by offering context-specific data. This adds to the conversation on telemedicine's potential to reduce the burden of chronic illness and enhance the health of the elderly, especially in areas where access to conventional healthcare services is scarce. The study's overall demonstrate how telemedicine may improve the treatment of chronic illnesses and the way senior citizens get healthcare, which will further our knowledge of the technology's wider effects on healthcare outcomes.

Effectiveness of Telemedicine Services in Enhancing Healthcare Access

The study's conclusions offer convincing information on how well telemedicine services might improve elderly patients' access to healthcare in isolated Saudi Arabian locales. The research clarifies the important role of telemedicine in removing geographical barriers and enhancing healthcare access through an investigation of multiple access indicators, including geographic location, demographic factors, and healthcare consumption patterns. Telemedicine treatments efficiently bridge the gap between senior people and healthcare practitioners, especially in areas with limited access to traditional healthcare facilities, by facilitating remote consultations, virtual follow-ups, and telemonitoring. The aforementioned results enhance our comprehension of the influence of telemedicine on healthcare accessibility and underscore its capability to mitigate healthcare inequalities among senior people residing in remote areas.

Clinical Outcomes Associated with Telemedicine Interventions

The study research on the therapeutic results of telemedicine interventions for senior citizens living in isolated Saudi Arabian areas. Through the assessment of chronic condition monitoring, medication adherence rates, and disease management ratings, the study offers important new information on how well telemedicine works to improve health outcomes for the elderly. The study's conclusions show the revolutionary potential of telemedicine in improving healthcare delivery by demonstrating beneficial clinical results, such as increased medication adherence, proactive monitoring of chronic illnesses, and enhanced disease management. These findings highlight the importance of telemedicine in raising the standard of care for elderly patients in remote locations and add to the increasing body of research demonstrating its ability to improve clinical outcomes.

Impact of Telemedicine Services on Patient Satisfaction and Cost-Effectiveness

The research investigates the effects of telemedicine services on patient satisfaction and cost-effectiveness when it comes to senior healthcare in Saudi Arabia's outlying districts. The study assesses patient perceptions, satisfaction levels, and the financial effects of telemedicine treatments using questionnaires, interviews, and data on healthcare usage. The results show that patients are highly satisfied with telemedicine services, which may be ascribed to elements like ease of use, accessibility, and high-quality medical treatment. The study also examines how cost-effective telemedicine is in comparison to conventional healthcare delivery techniques, taking into account things like lower travel costs, fewer hospital stays, and better resource allocation. These

discoveries advance our knowledge of telemedicine's effects on healthcare costs and patient happiness, which will guide future resource allocation plans and policy choices.

CONCLUSION

The usefulness of telemedicine services in meeting the healthcare requirements of senior patients living in isolated regions of Saudi Arabia has been clarified by this study. By means of an extensive examination of healthcare access, clinical results, patient contentment, and financial efficiency, the study has yielded significant knowledge regarding the revolutionary possibilities of telemedicine in enhancing healthcare provision for this particular population. The results underscore the noteworthy contribution of telemedicine towards surmounting geographical constraints, augmenting illness supervision, strengthening medication compliance, and elevating patient contentment. The study also highlights how affordable telemedicine is in comparison to conventional healthcare delivery techniques, highlighting its potential to produce long-term medical results.

Recommendations

The development and growth of telemedicine services should be a top priority for Saudi Arabia's healthcare authorities, especially in rural areas where access to medical facilities is restricted. This entails making investments in the infrastructure needed for communications, educating medical personnel about telemedicine, and educating senior patients about the advantages of telemedicine. Integration of Telemedicine into Healthcare Policies: To guarantee its broad acceptance and sustainability, telemedicine should be incorporated into national healthcare policies and plans. To ensure the quality, safety, and privacy of healthcare delivery, policymakers should work in conjunction with stakeholders, technology developers, and healthcare providers to create regulatory frameworks and standards for the deployment of telemedicine. Continuous Monitoring and Assessment: To determine the efficacy of telemedicine initiatives, pinpoint areas in need of development, and handle new issues, ongoing monitoring and assessment are crucial. In order to track important performance metrics, clinical results, patient satisfaction, and the cost-effectiveness of telemedicine services, healthcare institutions should set up strong monitoring and evaluation processes. Research and Innovation: To improve telemedicine technology and therapies' efficacy and meet changing healthcare demands, more research and innovation are required. To create cutting-edge telemedicine solutions that are specifically suited to the needs of senior populations living in distant places, researchers, academic institutions, and industry partners should work together on multidisciplinary research initiatives. Community Empowerment and involvement: Successful telemedicine program implementation depends on community empowerment and involvement. It is recommended that healthcare practitioners establish connections with patient advocacy organizations, local leaders, and communities to foster trust, increase awareness, and overcome social and cultural obstacles to the use of telemedicine.

REFERENCES

1. Al Saffer, Q., Al-Ghaith, T., Alshehri, A., Al-Mohammed, R., Al Homidi, S., Hamza, M. M., ... & Alazemi, N. (2021). The capacity of primary health care facilities in Saudi Arabia: infrastructure, services, drug availability, and human resources. *BMC health services research*, 21, 1-15. <https://doi.org/10.1186/s12913-021-06355-x>
2. Alaboudi, A., Atkins, A., Sharp, B., Balkhair, A., Alzahrani, M., & Sunbul, T. (2016). Barriers and challenges in adopting Saudi telemedicine network: The perceptions of decision makers of healthcare facilities in Saudi Arabia. *Journal of infection and public health*, 9(6), 725-733. <https://doi.org/10.1016/j.jiph.2016.09.001>
3. Alharthi, H., Alsubhi, M., Shamsan, A., & Househ, M. (2020). Challenges and opportunities of telemedicine services in Saudi Arabia: A qualitative study. *Journal of Infection and Public Health*, 13(4), 552-557.
4. Alhomoud, F., Dhillon, S., Aslanpour, Z., & Smith, F. (2015). South Asian and Middle Eastern patients' perspectives on medicine-related problems in the United Kingdom. *International journal of clinical pharmacy*, 37, 607-615. <https://doi.org/10.1007/s11096-015-0103-6>
5. Almalki, M., FitzGerald, G., & Clark, M. (2011). Health care system in Saudi Arabia: an overview. *EMHJ-Eastern Mediterranean Health Journal*, 17 (10), 784-793, 2011.
6. Bashshur, R. L., Shannon, G. W., Bashshur, N., & Yellowlees, P. M. (2016). The empirical evidence for telemedicine interventions in mental disorders. *Telemedicine and e-Health*, 22(2), 87-113. <https://doi.org/10.1089/tmj.2015.0206>

7. Bradford, N. K., Caffery, L. J., & Smith, A. C. (2015). Awareness, experiences and perceptions of telehealth in a rural Queensland community. *BMC health services research*, 15, 1-10. <https://doi.org/10.1186/s12913-015-1094-7>
8. Chandwani, R., De, R., & Dwivedi, Y. K. (2018). Telemedicine for low resource settings: Exploring the generative mechanisms. *Technological forecasting and social change*, 127, 177-187. <https://doi.org/10.1016/j.techfore.2017.06.014>
9. Cimperman, M., Brenčič, M. M., Trkman, P., & Stanonik, M. D. L. (2013). Older adults' perceptions of home telehealth services. *Telemedicine and e-Health*, 19(10), 786-790. <https://doi.org/10.1089/tmj.2012.0272>
10. Flodgren, G., Rachas, A., Farmer, A. J., Inzitari, M., & Shepperd, S. (2015). Interactive telemedicine: Effects on professional practice and health care outcomes. *Cochrane Database of Systematic Reviews*, (9).
11. General Authority for Statistics. (2020). Demographic Survey.
12. Hajesmaeel-Gohari, S., & Bahaadinbeigy, K. (2021). The most used questionnaires for evaluating telemedicine services. *BMC medical informatics and decision making*, 21(1), 1-11. <https://doi.org/10.1186/s12911-021-01407-y>
13. Hazlett-Stevens, H., Singer, J., & Chong, A. (2019). Mindfulness-based stress reduction and mindfulness-based cognitive therapy with older adults: A qualitative review of randomized controlled outcome research. *Clinical gerontologist*, 42(4), 347-358. <https://doi.org/10.1080/07317115.2018.1518282>
14. Mayberry, R. M., Nicewander, D. A., Qin, H., & Ballard, D. J. (2006, April). Improving quality and reducing inequities: a challenge in achieving best care. In *Baylor University Medical Center Proceedings* (Vol. 19, No. 2, pp. 103-118). Taylor & Francis. <https://doi.org/10.1080/08998280.2006.11928138>
15. Nelson, H. D. (2024). Systematic reviews to answer health care questions. Lippincott Williams & Wilkins.
16. Omboni, S., Padwal, R. S., Alessa, T., Benczúr, B., Green, B. B., Hubbard, I., ... & Wang, J. (2022). The worldwide impact of telemedicine during COVID-19: current evidence and recommendations for the future. *Connected health*, 1, 7. <https://doi.org/10.20517/2Fch.2021.03>
17. Paul, D. L., Pearlson, K. E., & McDaniel, R. R. (1999). Assessing technological barriers to telemedicine: technology-management implications. *IEEE Transactions on engineering management*, 46(3), 279-288. <https://doi.org/10.1109/17.775280>
18. Rachas, A., Farmer, A. J., Inzitari, M., & Shepperd, S. (2015). Interactive telemedicine: effects on professional practice and health care outcomes. *Cochrane Database of Systematic Reviews*, (9). <https://doi.org/10.1002/14651858.CD002098.pub2>
19. Rosli, M. S., Saleh, N. S., Md. Ali, A., Abu Bakar, S., & Mohd Tahir, L. (2022). A Systematic review of the technology acceptance model for the sustainability of higher education during the COVID-19 pandemic and identified research gaps. *Sustainability*, 14(18), 11389. <https://doi.org/10.3390/su141811389>
20. Salam, A. A. (2023). Ageing in Saudi Arabia: new dimensions and intervention strategies. *Scientific Reports*, 13(1), 4035. <https://doi.org/10.1038/s41598-022-25639-8>
21. Tannous, W. K., & Quilty, K. (2022). Ageing: Everybody's Doing It—Life-Affirming Technology and the Longevity Economy. In *Research Anthology on Supporting Healthy Aging in a Digital Society* (pp. 556-570). IGI Global. <https://doi.org/10.4018/978-1-6684-5295-0.ch031>
22. United Nations. (2019). World population prospects 2019: Highlights. United Nations Publications.
23. Van den Berg, N., Schumann, M., Kraft, K., & Hoffmann, W. (2012). Telemedicine and telecare for older patients—a systematic review. *Maturitas*, 73(2), 94-114. <https://doi.org/10.1016/j.maturitas.2012.06.010>
24. World Health Organization. (2010). Telemedicine: opportunities and developments in member states. Report on the second global survey on eHealth. World Health Organization. <http://apps.who.int/iris/handle/10665/44497>
25. World Health Organization. (2015). World report on ageing and health. World Health Organization.