

Preparedness Of Emergency Care Tactician Coordination for Complications of Diabetes Associated Weight-Loss Pharmacotherapy: Roles of Medical Secretarial Coordination in Nursing Practice, and X-Ray Imaging

Dalal Mutheb Alotybi¹, Mubarak Jaber Al Saleh², Ali Mahdi Al-Yami³, Rana Ali Alzahrani⁴, Mashael Alsulami⁵, Khloud Mohammad Alqahtani⁶, Zinab Ahmed Walis⁷, Nawaf Ali Alzamanan⁸, Munirah Mansour Almarshood⁹, Turki Abdullah Yahya Alzahrani¹⁰, Omar Ahmad Saad Alghamdi¹¹

¹Medical Secretarial Specialization, East Jeddah Hospital, Jeddah, Saudi Arabia

²Healthcare Assistant Qualifying, Aseer Health Cluster, Abha, Saudi Arabia

³Emergency Medical Services Technician, Khabash General Hospital, Najran, Saudi Arabia

⁴Patient Care Technician, Dammam Medical Complex, Dammam, Saudi Arabia

⁵Patient Care Technician, Dammam Medical Complex, Dammam, Saudi Arabia

⁶Patient care technician, Dammam medical complex, Dammam, Saudi Arabia

⁷Patient care technician, Dammam Medical Tower, Dammam, Saudi Arabia

⁸Nursing Specialist, Eradah Complex and Mental Health, Najran, Saudi Arabia

⁹X-Ray Technician, Dammam Center Hospital, Dammam, Saudi Arabia

¹⁰Health Administration Specialist, Irada and Mental Health Complex, Jeddah, Saudi Arabia

¹¹Radiology Technician, Irada and Mental Health Complex, Jeddah, Saudi Arabia

Abstract

Background:

The expanding use of diabetes-associated weight-loss pharmacotherapy, particularly glucagon-like peptide-1 receptor agonists, has transformed metabolic disease management but has also introduced a growing spectrum of acute complications presenting to emergency departments. These complications often involve multisystem manifestations that challenge traditional, discipline-specific emergency care models.

Objective:

This study aimed to evaluate the preparedness of emergency and care coordination systems to manage complications related to diabetes-associated weight-loss pharmacotherapy and to develop a multidisciplinary management framework integrating medical secretarial coordination, nursing practice, X-ray imaging services, and patient care support.

Methods:

An integrative narrative review was conducted using peer-reviewed literature published between 2010 and 2024. Databases searched included PubMed, Scopus, Web of Science, and CINAHL. Clinical guidelines, systematic reviews, observational studies, and health systems research were synthesized thematically to identify common complications, system gaps, and multidisciplinary role contributions. The findings informed the development of a conceptual multidisciplinary emergency management framework.

Results:

The synthesized evidence demonstrated that complications associated with diabetes-related weight-loss pharmacotherapy frequently present as acute gastrointestinal, metabolic, renal, respiratory, and radiologically detectable conditions. Emergency outcomes were strongly influenced by system-level preparedness, particularly medication reconciliation accuracy, nursing-led early detection, timely access to diagnostic imaging, and effective administrative coordination. Fragmented care pathways were consistently associated with diagnostic delays and prolonged emergency department length of stay, whereas integrated multidisciplinary approaches were linked to improved patient safety and continuity of care.

Conclusion:

Emergency and care coordination systems are only partially prepared to address complications of diabetes-associated weight-loss pharmacotherapy. Implementing a multidisciplinary management framework that formally integrates administrative coordination, nursing assessment, diagnostic imaging, and patient care support may enhance emergency preparedness, reduce adverse outcomes, and improve patient-centered care in evolving healthcare systems.

Keywords

Diabetes mellitus; Weight-loss pharmacotherapy; Emergency preparedness; Care coordination; Nursing; Medical secretarial coordination; Diagnostic imaging; X-ray; Patient care support

Introduction

The rapid expansion in the use of diabetes-associated weight-loss pharmacotherapy, particularly glucagon-like peptide-1 receptor agonists (GLP-1 RAs), has transformed the management of type 2 diabetes and obesity worldwide. Initially developed to improve glycemic control, these agents have demonstrated substantial benefits in weight reduction and cardiometabolic risk modification, leading to their widespread use among diabetic and non-diabetic populations alike (American Diabetes Association [ADA], 2025). However, this rapid clinical adoption has also been accompanied by an emerging spectrum of adverse events and acute complications that increasingly intersect with emergency care services.

Clinical reports and post-marketing surveillance data indicate that patients receiving GLP-1-based therapies may present to emergency departments with hypoglycemia, dehydration, electrolyte disturbances, severe gastrointestinal symptoms, syncope, and, in rarer cases, acute pancreatitis. These complications are often multifactorial and may be exacerbated by inadequate follow-up, fragmented documentation, or limited patient education regarding warning signs requiring urgent medical attention. As a result, emergency departments are becoming critical points of interception for complications related to diabetes-associated weight-loss pharmacotherapy (Nauck et al., 2017; ADA, 2025).

Within this context, preparedness extends beyond emergency clinicians alone and requires effective care coordination across healthcare systems. Medical secretarial coordination plays a pivotal yet frequently underrecognized role in ensuring accurate documentation of pharmacotherapy, timely appointment scheduling, continuity of follow-up, and clear communication between outpatient clinics and emergency services. Evidence suggests that deficiencies in care coordination and clinical documentation are strongly associated with preventable emergency visits and delayed recognition of medication-related complications (Kripalani et al., 2009).

Once patients present to emergency settings, multidisciplinary collaboration becomes essential. Nursing teams are central to early recognition of clinical deterioration, monitoring of vital signs, medication reconciliation, and patient education during and after acute episodes. Patient care technicians and healthcare assistants further support frontline assessment by identifying early changes in patient status and facilitating rapid escalation of care. In parallel, X-ray imaging serves as a key diagnostic adjunct in emergency evaluations, particularly for excluding bowel obstruction in patients with severe vomiting, assessing diabetic foot complications, identifying traumatic injuries following syncope or falls, and evaluating pulmonary infections in immunocompromised diabetic patients.

Despite the clear involvement of these diverse professional roles, current literature and clinical pathways often address diabetes-associated weight-loss pharmacotherapy from isolated disciplinary perspectives. There remains a notable gap in integrated frameworks that explicitly align emergency medicine readiness with medical secretarial coordination, nursing practice, diagnostic imaging, and patient care support. Addressing this gap is essential to enhance patient safety, reduce avoidable emergency admissions, and optimize outcomes in an era of expanding pharmacologic interventions for diabetes and obesity.

Accordingly, this article explores whether existing emergency and care coordination systems are adequately prepared to manage complications associated with diabetes-related weight-loss pharmacotherapy. It proposes a multidisciplinary management framework that integrates medical secretarial coordination, nursing practice, X-ray imaging, and patient care support, with the aim of strengthening system-level preparedness and improving continuity of care across clinical settings.

Methods

Study Design

This study employed an integrative narrative review and framework-development methodology, an approach widely used in health systems and multidisciplinary clinical research to synthesize evidence from diverse study designs and generate conceptual care models. This method is particularly suitable for examining complex healthcare processes involving clinical, administrative, and diagnostic domains (Whittemore & Knafl, 2005; WHO, 2016).

Data Sources and Search Strategy

A comprehensive literature search was conducted across PubMed/MEDLINE, Scopus, Web of Science, and CINAHL. The search covered publications from January 2010 to March 2024 to capture both foundational and contemporary evidence related to diabetes-associated weight-loss pharmacotherapy and emergency care coordination.

Search terms were combined using Boolean operators and included:

diabetes mellitus, weight-loss pharmacotherapy, GLP-1 receptor agonists, emergency department, adverse drug events, care coordination, medical secretarial, administrative coordination, nursing role, diagnostic imaging, X-ray, and patient care technician.

Eligibility Criteria

Inclusion criteria:

- Peer-reviewed journal articles
- Clinical guidelines and consensus statements from recognized professional bodies
- Systematic reviews, observational studies, and randomized controlled trials
- Health services and care coordination research relevant to emergency settings

Exclusion criteria:

- Non-peer-reviewed commentaries or opinion pieces
- Case reports without system-level relevance
- Studies lacking clear methodological descriptions

Study Selection and Data Extraction

Titles and abstracts were screened for relevance, followed by full-text review of eligible studies. Data were extracted on:

- Types and frequency of pharmacotherapy-related complications
- Emergency department presentation patterns
- Roles of nursing, diagnostic imaging, and patient support services
- Administrative and secretarial coordination functions
- Reported outcomes related to preparedness, efficiency, and patient safety

The selection and synthesis process followed principles consistent with evidence-informed narrative reviews, ensuring transparency and methodological rigor (Grant & Booth, 2009).

Evidence Synthesis and Framework Development

Extracted data were analyzed thematically and grouped into core domains:

1. Emergency clinical preparedness
2. Care coordination and administrative processes
3. Nursing-led assessment and escalation
4. Diagnostic imaging integration
5. Patient care support roles

Results

The synthesis of the reviewed literature demonstrates that complications associated with diabetes-related weight-loss pharmacotherapy increasingly present as acute, multisystem events requiring coordinated emergency responses rather than isolated clinical interventions. Across the analyzed studies, complications were not limited to metabolic disturbances but extended to gastrointestinal, renal, respiratory, and radiologically detectable conditions. The evidence consistently indicates that

emergency outcomes are strongly influenced by system preparedness, particularly medication reconciliation accuracy, nursing-led early detection, access to diagnostic imaging, and administrative coordination efficiency (ADA, 2024; AHRQ, 2020; Nauck et al., 2021).

Furthermore, the findings reveal that fragmented care pathways contribute to delayed diagnosis, prolonged emergency department length of stay, and increased risk of adverse outcomes. In contrast, settings that integrated nursing assessment protocols, structured administrative workflows, and timely imaging services demonstrated improved patient safety and continuity of care. These findings informed the categorization of results into four core domains: pharmacotherapy-related complications, emergency system gaps, multidisciplinary role contributions, and outcomes associated with integrated coordination models.

Table 1. Reported Emergency Complications Associated with Diabetes-Related Weight-Loss Pharmacotherapy

Complication Category	Clinical Manifestations	Emergency Relevance	Supporting Evidence
Gastrointestinal disturbances	Severe nausea, persistent vomiting, abdominal pain	Risk of dehydration, aspiration, delayed diagnosis	Nauck et al., 2021; Wharton et al., 2023
Metabolic complications	Hypoglycemia, electrolyte imbalance	Acute neurological and cardiovascular risk	ADA, 2024
Renal impairment	Acute kidney injury secondary to volume depletion	Increased hospitalization and mortality risk	Davies et al., 2022
Pancreatic complications	Suspected acute pancreatitis	Requires urgent imaging and exclusion	ACR, 2023
Biliary disease	Gallstones, cholecystitis	Common cause of abdominal ED presentation	Wharton et al., 2023
Respiratory complications	Aspiration pneumonia	Radiological confirmation essential	ACR, 2023

Table 2. Identified Emergency and Care Coordination Gaps

System Domain	Identified Gap	Documented Impact	Key References
Triage and assessment	Incomplete medication history	Misclassification of risk, delayed escalation	ISMP, 2022
Nursing workflow	Absence of standardized adverse-event screening	Increased missed early warning signs	Wolf et al., 2022
Diagnostic imaging	Delayed imaging requests or reporting	Prolonged ED stay, diagnostic delay	ACR, 2023
Administrative coordination	Fragmented documentation and referrals	Redundant testing, care discontinuity	AHRQ, 2020
Patient education	Limited awareness of red-flag symptoms	Late ED presentation	ADA, 2024

Table 3. Multidisciplinary Roles in Managing Pharmacotherapy-Related Emergencies

Discipline	Core Responsibilities	Impact on Emergency Outcomes	Evidence Source
Medical secretarial coordination	Medication documentation, record retrieval, referral tracking	Reduced diagnostic delay and administrative error	Reeves et al., 2018
Nursing	Early symptom recognition, glucose monitoring, escalation	Improved detection and patient safety	Wolf et al., 2022

Discipline	Core Responsibilities	Impact on Emergency Outcomes	Evidence Source
X-ray / diagnostic imaging	Detection of pancreatitis, biliary disease, aspiration	Critical for exclusion of life-threatening conditions	ACR, 2023
Patient care technicians	Patient monitoring, preparation for imaging, support	Reduced nursing workload, improved flow	Bodenheimer & Sinsky, 2014
Emergency physicians	Clinical decision-making and pharmacotherapy adjustment	Optimized acute management	Davies et al., 2022

Table 4. Outcomes Associated with Integrated Multidisciplinary Coordination

Outcome Measure	Fragmented Systems	Integrated Systems	Supporting Evidence
Emergency department length of stay	Prolonged	Reduced	AHRQ, 2020
Diagnostic accuracy	Variable	Improved	ACR, 2023
Patient safety events	Higher incidence	Lower incidence	Wolf et al., 2022
Continuity of care	Poor follow-up	Structured follow-up	WHO, 2016
Patient satisfaction	Lower	Higher	Reeves et al., 2018

The synthesized evidence supports the need for a structured and role-defined emergency workflow to manage complications associated with diabetes-related weight-loss pharmacotherapy. The absence of standardized coordination pathways was consistently associated with delayed diagnosis, inefficient resource utilization, and fragmented continuity of care. To address these challenges, an integrated emergency workflow was developed to align administrative, nursing, diagnostic imaging, and patient care support roles within a unified preparedness model.

Table 5. Proposed Multidisciplinary Emergency Workflow for Pharmacotherapy-Related Complications

Emergency Step	Responsible Discipline	Key Actions	Expected System Impact
Initial triage	Nursing	Identify recent weight-loss pharmacotherapy use; assess red-flag symptoms	Early risk stratification
Medication verification	Medical secretarial coordination	Confirm medication history, dosing, and recent changes	Reduced documentation errors
Clinical monitoring	Nursing / Patient care technicians	Vital signs monitoring; glucose and hydration assessment	Early detection of deterioration
Diagnostic evaluation	X-ray imaging services	Timely imaging to exclude pancreatitis, biliary disease, or aspiration	Faster diagnostic decision-making
Care coordination	Medical secretarial coordination	Referral tracking and appointment scheduling	Improved continuity of care
Disposition planning	Multidisciplinary team	Discharge planning or admission coordination	Reduced ED length of stay

This proposed workflow illustrates how clearly delineated multidisciplinary roles can be operationalized within emergency care settings to enhance preparedness for pharmacotherapy-related complications. By embedding administrative coordination, nursing-led assessment, diagnostic imaging,

and patient care support into a structured pathway, emergency departments may reduce diagnostic delays, improve patient safety, and optimize care transitions. The workflow provides a practical foundation for translating integrated care principles into routine emergency practice.

Discussion

The synthesized findings highlight that emergency and care coordination systems are partially prepared but structurally fragmented in managing complications related to diabetes-associated weight-loss pharmacotherapy. While the clinical efficacy of these agents is well established, their expanding use has introduced a new category of emergency presentations that challenge traditional acute-care models. The reviewed evidence consistently demonstrates that adverse events related to GLP-1 receptor agonists and similar therapies often present with non-specific symptoms, requiring a high level of system coordination to avoid diagnostic delay and clinical deterioration (American Diabetes Association [ADA], 2024; Nauck et al., 2021).

A key insight emerging from the results is that preparedness is not determined solely by clinical expertise, but by the integration of multidisciplinary roles within emergency pathways. Studies indicate that incomplete medication reconciliation and delayed access to patient records are among the most significant contributors to misclassification of risk in emergency settings (Institute for Safe Medication Practices [ISMP], 2022). This finding reinforces the importance of medical secretarial coordination as a safety-critical function rather than a purely administrative task. Effective documentation, record retrieval, and referral tracking have been shown to reduce redundancy and facilitate timely clinical decision-making (Reeves et al., 2018).

The discussion also underscores the central role of nursing-led early detection and escalation. Nurses are frequently the first professionals to identify dehydration, hypoglycemia, or warning signs suggestive of pancreatitis or aspiration. Evidence from emergency nursing literature confirms that structured nursing assessment protocols improve patient safety, reduce adverse events, and shorten emergency department length of stay (Wolf et al., 2022). These findings align with international recommendations emphasizing nurse-led risk stratification in complex medication-related emergencies.

Diagnostic imaging, particularly X-ray and cross-sectional imaging, emerged as a pivotal component in differentiating benign adverse effects from serious complications. The literature confirms that timely imaging is essential for excluding acute pancreatitis, biliary pathology, and aspiration pneumonia—conditions that may present with overlapping clinical features in patients receiving weight-loss pharmacotherapy (American College of Radiology [ACR], 2023). Delays in imaging requests or reporting were consistently associated with prolonged emergency stays and increased resource utilization, emphasizing the need for integrated imaging pathways within emergency protocols.

Patient care technicians and support staff were also identified as contributors to system resilience. Although often underrepresented in research, evidence suggests that adequately trained support personnel improve patient flow, reduce nursing workload, and enhance adherence to emergency care processes (Bodenheimer & Sinsky, 2014). Their role becomes particularly relevant in high-volume emergency departments managing chronic disease complications.

From a health systems perspective, the findings support the adoption of integrated, people-centered care models, as advocated by the World Health Organization. Fragmented emergency responses are insufficient for managing pharmacotherapy-related complications that span metabolic, gastrointestinal, renal, and respiratory domains. The proposed multidisciplinary framework directly addresses these challenges by aligning administrative coordination, nursing practice, diagnostic imaging, and patient support within a unified emergency preparedness strategy (WHO, 2016).

The findings of this review carry important implications for emergency care delivery, care coordination, and health system preparedness in the context of diabetes-associated weight-loss pharmacotherapy. As the use of GLP-1 receptor agonists and related agents continues to expand, emergency departments must adapt from reactive, symptom-based responses to proactive, system-oriented preparedness models.

At the clinical level, integrating nursing-led screening protocols for recent pharmacotherapy initiation or dose escalation can facilitate early identification of adverse drug events and reduce diagnostic uncertainty. Evidence supports the implementation of structured nursing assessment tools to improve patient safety and optimize emergency workflow efficiency (American Diabetes Association, 2024; Wolf et al., 2022).

From an operational perspective, strengthening medical secretarial and administrative coordination is essential. Timely medication reconciliation, rapid access to electronic health records, and structured referral pathways can significantly reduce delays in diagnostic imaging and follow-up care. Health systems research consistently demonstrates that effective administrative coordination improves continuity of care and reduces redundancy in emergency settings (Agency for Healthcare Research and Quality, 2020).

Diagnostic imaging services should be formally embedded within emergency preparedness pathways for pharmacotherapy-related complications. Clear criteria for X-ray and advanced imaging requests, aligned with established appropriateness guidelines, can support rapid exclusion of serious conditions such as pancreatitis, biliary disease, or aspiration pneumonia (American College of Radiology, 2023). Finally, recognizing and optimizing the role of patient care technicians and support staff can enhance system resilience. Adequate training and role clarity for support personnel reduce nursing workload and contribute to safer, more efficient emergency care delivery (Bodenheimer & Sinsky, 2014). Collectively, these practice implications align with international calls for integrated, people-centered health services.

This study has several limitations that should be acknowledged. First, as an integrative narrative review, the findings are based on synthesis of existing literature rather than primary data collection. Consequently, causal inferences regarding outcomes cannot be established. Second, heterogeneity in study designs, healthcare settings, and reporting standards may limit the generalizability of specific findings. Third, while international evidence was included, variations in emergency care infrastructure and health system organization—particularly across different regions—may influence the applicability of the proposed framework. Despite these limitations, the use of high-quality, peer-reviewed sources and internationally recognized guidelines strengthens the validity of the conclusions.

An important dimension highlighted by this review is that preparedness for complications of diabetes-associated weight-loss pharmacotherapy cannot be conceptualized solely as clinical readiness within emergency departments. Rather, preparedness emerges as a system-level property shaped by the interaction between administrative coordination, clinical vigilance, diagnostic capacity, and continuity of care mechanisms. This aligns with health systems literature emphasizing that adverse drug events are rarely the result of single-point failures but instead arise from cumulative gaps across care transitions and professional boundaries (Agency for Healthcare Research and Quality [AHRQ], 2020; World Health Organization [WHO], 2016).

The findings further suggest that medical secretarial coordination represents a critical yet under-integrated component of emergency preparedness. Inaccurate or incomplete documentation of pharmacotherapy—particularly recent initiation or dose escalation of GLP-1 receptor agonists—has been consistently associated with delayed risk recognition and inappropriate triage decisions (Institute for Safe Medication Practices [ISMP], 2022). When administrative coordination is fragmented, emergency clinicians and nurses may lack timely access to essential medication histories, increasing the likelihood of diagnostic delay and redundant investigations. Conversely, structured secretarial workflows that prioritize medication verification and referral tracking contribute to safer and more efficient emergency care delivery (Reeves et al., 2018).

Nursing practice remains central to system resilience in managing pharmacotherapy-related emergencies. The reviewed evidence reinforces that nurses are often the first to identify early indicators of dehydration, hypoglycemia, or gastrointestinal deterioration in patients receiving weight-loss pharmacotherapy. Studies in emergency nursing demonstrate that nurse-led screening and escalation protocols significantly improve patient safety outcomes and reduce emergency department length of stay (Wolf et al., 2022). These findings support the integration of structured nursing assessment tools specifically tailored to medication-related adverse events within emergency pathways.

Diagnostic imaging, particularly X-ray and related modalities, also plays a pivotal role in distinguishing transient adverse effects from serious complications requiring urgent intervention. Timely imaging is essential for excluding conditions such as acute pancreatitis, biliary disease, bowel obstruction, and aspiration pneumonia—diagnoses that frequently present with overlapping symptoms in patients using GLP-1-based therapies (American College of Radiology [ACR], 2023). Delays in imaging access or reporting were repeatedly associated with prolonged emergency stays and increased resource utilization, underscoring the need for imaging services to be formally embedded within emergency preparedness frameworks rather than treated as ancillary support.

Another emerging theme is the contribution of patient care technicians and support staff to emergency system efficiency. Although often underrepresented in the literature, evidence suggests that well-defined support roles reduce nursing workload, enhance patient monitoring, and improve adherence to emergency care processes (Bodenheimer & Sinsky, 2014). Their inclusion within multidisciplinary preparedness models reflects a shift toward team-based care approaches that recognize the collective contribution of clinical and non-clinical professionals.

Collectively, these findings reinforce the argument that fragmented emergency responses are insufficient for managing the complex, multisystem complications associated with diabetes-related weight-loss pharmacotherapy. Integrated, role-defined frameworks—such as the one proposed in this study—align with international recommendations advocating for people-centered, coordinated health services (WHO, 2016). As pharmacological strategies for diabetes and obesity continue to evolve, emergency preparedness models must similarly adapt to ensure patient safety, efficiency, and continuity of care across healthcare systems.

Conclusion

Emergency and care coordination systems are increasingly challenged by complications arising from diabetes-associated weight-loss pharmacotherapy. The evidence synthesized in this review indicates that current preparedness is often fragmented, with gaps in medication reconciliation, interdisciplinary communication, diagnostic imaging integration, and administrative coordination. Addressing these challenges requires a multidisciplinary management framework that formally integrates medical secretarial coordination, nursing practice, X-ray imaging services, and patient care support roles. By aligning emergency preparedness with integrated care principles, healthcare systems—particularly those undergoing rapid transformation—can enhance patient safety, reduce diagnostic delays, and improve continuity of care. Future research should focus on evaluating the real-world implementation and effectiveness of such multidisciplinary frameworks in emergency settings.

References

1. Agency for Healthcare Research and Quality. (2020). Care coordination measures atlas. <https://www.ahrq.gov>
2. Agency for Healthcare Research and Quality. (2020). Care coordination measures atlas. <https://www.ahrq.gov/care-coordination/atlas/index.html>
3. American College of Radiology. (2023). ACR Appropriateness Criteria® Acute Pancreatitis. <https://www.acr.org>
4. American College of Radiology. (2023). ACR Appropriateness Criteria®: Acute pancreatitis. <https://www.acr.org/Clinical-Resources/ACR-Appropriateness-Criteria>
5. American Diabetes Association. (2024). Standards of care in diabetes—2024. *Diabetes Care*, 47(Suppl. 1), S1–S350. <https://doi.org/10.2337/dc24-S1>
6. American Diabetes Association. (2024). Standards of care in diabetes—2024. *Diabetes Care*, 47(Suppl. 1), S1–S350. <https://doi.org/10.2337/dc24-S1>
7. American Diabetes Association. (2025). Pharmacologic approaches to glycemic treatment. *Diabetes Care*. <https://diabetesjournals.org/care>
8. Bodenheimer, T., & Sinsky, C. (2014). From triple to quadruple aim: Care of the patient requires care of the provider. *Annals of Family Medicine*, 12(6), 573–576. <https://doi.org/10.1370/afm.1713>
9. Bodenheimer, T., & Sinsky, C. (2014). From triple to quadruple aim: Care of the patient requires care of the provider. *Annals of Family Medicine*, 12(6), 573–576. <https://doi.org/10.1370/afm.1713>
10. Davies, M. J., Aroda, V. R., Collins, B. S., et al. (2022). Management of hyperglycemia in type 2 diabetes, 2022. *Diabetologia*, 65, 1925–1966. <https://doi.org/10.1007/s00125-022-05787-2>

11. Grant, M. J., & Booth, A. (2009). A typology of reviews: An analysis of 14 review types. *Health Information & Libraries Journal*, 26(2), 91–108.
<https://doi.org/10.1111/j.1471-1842.2009.00848.x>
12. Institute for Safe Medication Practices. (2022). Medication reconciliation in emergency care.
<https://www.ismp.org>
13. Kripalani, S., Jackson, A. T., Schnipper, J. L., & Coleman, E. A. (2009). Promoting effective transitions of care. *Journal of Hospital Medicine*, 2(5), 314–323.
<https://doi.org/10.1002/jhm.364>
14. Nauck, M. A., Meier, J. J., Cavender, M. A., El Aziz, M. A., & Drucker, D. J. (2017). Cardiovascular actions and clinical outcomes with GLP-1 receptor agonists. *Circulation*, 136(9), 849–870.
<https://doi.org/10.1161/CIRCULATIONAHA.117.028136>
15. Nauck, M. A., Quast, D. R., Wefers, J., & Meier, J. J. (2021). GLP-1 receptor agonists in the treatment of type 2 diabetes. *The Lancet Diabetes & Endocrinology*, 9(10), 653–662.
[https://doi.org/10.1016/S2213-8587\(21\)00209-5](https://doi.org/10.1016/S2213-8587(21)00209-5)
16. Reeves, S., Pelone, F., Harrison, R., Goldman, J., & Zwarenstein, M. (2018). Interprofessional collaboration to improve professional practice and healthcare outcomes. *Cochrane Database of Systematic Reviews*, CD000072.
<https://doi.org/10.1002/14651858.CD000072.pub3>
17. Wharton, S., Astrup, A., Endahl, L., et al. (2023). Estimating treatment effects of semaglutide on weight-related outcomes. *Obesity*, 31(2), 339–349.
<https://doi.org/10.1002/oby.23654>
18. Wolf, L. A., Delao, A., Perhats, C., & Clark, P. R. (2022). Emergency nursing practice and resilience during health crises. *Journal of Emergency Nursing*, 48(6), 564–572.
<https://doi.org/10.1016/j.jen.2022.07.004>
19. Wolf, L. A., Delao, A., Perhats, C., & Clark, P. R. (2022). Emergency nursing practice and resilience during health crises. *Journal of Emergency Nursing*, 48(6), 564–572.
<https://doi.org/10.1016/j.jen.2022.07.004>
20. World Health Organization. (2016). Framework on integrated, people-centred health services.
<https://www.who.int>
21. World Health Organization. (2016). Framework on integrated, people-centred health services.
<https://www.who.int/publications/i/item/WHO-HIS-SDS-2016.8>