

# Falls Prevention and Mobility Support in Geriatric Nursing: Evidence-Based Interventions

**Yousef Eisa Alrefai<sup>1</sup>, Maha Dhafer Ayedh Al-Omari<sup>2</sup>, Fatima Ahmed Abdullah Sabr<sup>3</sup>, Khaled Salman Ali Al-Jahni<sup>4</sup>, Nouf Mohammed Aqili Al-Anzi<sup>5</sup>, Haya Hamad Mohammed Al-Shahrani<sup>6</sup>, Nihaya Dhawi Al-Anzi<sup>7</sup>, Fatima Dafer Alamri<sup>8</sup>, Ashwaq Abdullah Al Sharif<sup>9</sup>, Zainat Abdullah Al Shamrani<sup>10</sup>, Mansoura Fayhan Ashwi Al Ruwaili<sup>11</sup>, Amna Omar Qaisi<sup>12</sup>, Samar Shuai Barakat<sup>13</sup>, Shuroq Awaidah Alrwaili<sup>14</sup>, Eid Awadh Atiyyah Alanazi<sup>15</sup>, Reem Jafar Bantan<sup>16</sup>, Layla Yahya Al Fahmi<sup>17</sup>**

<sup>1</sup>. Nursing Technician, Executive Administration for Providing Health Care, Medina

<sup>2</sup>. Nursing Technician, Al-Quba Health Center, Bisha

<sup>3</sup>. Nursing Technician, Diabetes and Endocrinology Center, Bisha

<sup>4</sup>. Nursing, Madinah General Hospital, Madinah

<sup>5</sup>. Nursing Technician, East Al-Muhammadiyah Health Center, Hafar Al-Batin

<sup>6</sup>. Nursing, Public Security Training City in East Riyadh, Riyadh

<sup>7</sup>. Nursing, Al-Manar Health Center, Riyadh

<sup>8</sup>. Nursing technician, Ministry of Health Office, Bisha

<sup>9</sup>. Nursing Specialist, Al Suwaidi Western Health Center, Riyadh

<sup>10</sup>. Nursing Technician, Al Alia Primary Health Care Center, Medina

<sup>11</sup>. Nursing Technician, Hadeeb Health Center, Sakaka Al Jawf

<sup>12</sup>. Nursing Technician, Irada and Mental Health Complex (Irada Services), Jeddah

<sup>13</sup>. Nursing Technician, King Faisal Health Center

<sup>14</sup>. Nursing, Maternity and Children Hospital, Sakaka, Al-Jouf Region, Saudi Arabia (KSA)

<sup>15</sup>. Nursing Technician, Alazizah – Al-Jouf Health Cluster, Al-Qurayyat, Saudi Arabia (KSA)

<sup>16</sup>. Nursing Technician, King Saud Medical City, Riyadh, Saudi Arabia (KSA)

<sup>17</sup>. Nursing Technician, Al-Ajawid Primary Health Care Center (affiliated with Al-Thagr Hospital), Jeddah, Saudi Arabia (KSA)

## Abstract

Falls among older adults represent a critical public health concern, significantly contributing to morbidity and mortality in this population. This review article examines evidence-based interventions in geriatric nursing aimed at preventing falls and enhancing mobility among older adults. The multifactorial nature of falls necessitates a comprehensive understanding of both intrinsic and extrinsic risk factors, including age-related physiological changes, cognitive impairments, environmental hazards, and chronic health conditions. Geriatric nurses play a vital role in conducting thorough risk assessments, utilizing standardized tools to identify individuals at high risk for falls. The article highlights several key intervention strategies, including tailored exercise programs that focus on strength, balance, and flexibility, which have been shown to reduce fall risk significantly. Additionally, environmental modifications, such as improving home safety through the removal of hazards and the installation of assistive devices, are discussed as essential components of a fall prevention strategy. Medication management is also emphasized, as polypharmacy can increase fall risk; therefore, regular medication reviews and patient education are critical for optimizing pharmacotherapy. Furthermore, the importance of education and empowerment for both older adults and their caregivers is underscored, as knowledge about falls prevention can lead to proactive engagement in safety measures. Interdisciplinary collaboration is essential in creating comprehensive care plans that address the diverse needs of older adults, ensuring a holistic approach to falls prevention. Ultimately, this review underscores the significance of implementing evidence-based interventions in geriatric nursing to mitigate fall risks and promote mobility. By addressing the complex interplay of risk factors and fostering a supportive environment, healthcare professionals can enhance the quality of life for older adults, enabling them to maintain independence and reduce the fear associated with falling. The findings of this review advocate for a proactive, patient-centered approach to falls prevention in geriatric nursing practice.

## Introduction

Falls represent one of the most significant health threats to the elderly population, presenting a complex challenge that involves multifactorial risk factors and significant consequences. The World Health Organization (WHO) has identified falls as a leading cause of injury and mortality among

older adults, emphasizing the need for effective prevention strategies [1]. The statistics surrounding falls are alarming; it is estimated that one in three older adults experiences a fall each year, and among those who fall, approximately 20% to 30% sustain moderate to severe injuries, such as fractures or head trauma. These injuries not only lead to physical limitations but can also result in long-term disability, loss of independence, and even death. As the global population ages, with projections indicating that the number of individuals aged 65 and older will double by 2050, the incidence of falls is expected to rise significantly. This demographic shift underscores the urgent need for healthcare professionals, particularly geriatric nurses, to implement evidence-based interventions that address both the prevention of falls and the promotion of mobility among older adults [2].

Geriatric nursing encompasses a holistic approach to the care of older adults, focusing not only on the physical aspects of health but also on the psychological and social dimensions that contribute to overall well-being. This comprehensive perspective recognizes that the health of older adults is influenced by a myriad of factors, including their medical history, current health status, mental health, social support systems, and living environments. Within this framework, falls prevention and mobility support are critical components of care that can significantly enhance the quality of life for older adults. Falls can lead to a profound sense of fear and anxiety, which may deter older adults from engaging in physical activity, further exacerbating their risk of falls. This cycle of fear and inactivity can diminish their overall health and well-being, making it imperative for healthcare providers to address these issues proactively [3]. The complexity of falls among older adults necessitates a multifaceted approach to prevention. Geriatric nurses play a pivotal role in this effort, as they are often the primary healthcare providers for older adults in various settings, including hospitals, nursing homes, and community health programs [4]. Their unique position allows them to conduct comprehensive assessments, identify risk factors, and develop tailored interventions that address the specific needs of each patient. This review article explores the evidence-based interventions that geriatric nurses can employ to prevent falls and support mobility. It will delve into the underlying risk factors that contribute to falls, the importance of interdisciplinary collaboration in creating effective prevention strategies, and the role of education and empowerment in promoting safety and independence among older adults [5].

In addition to physical health considerations, it is essential to recognize the psychological impacts of falls on older adults. The fear of falling can lead to a decrease in physical activity, which, in turn, can result in further deconditioning and an increased risk of falls. This vicious cycle highlights the need for interventions that not only focus on physical safety but also address the emotional and psychological well-being of older adults. Strategies that promote confidence in mobility and encourage participation in physical activities can significantly mitigate the fear of falling. Therefore, geriatric nurses must be equipped with the knowledge and skills to address these psychological dimensions in their practice [6].

Moreover, the role of family members and caregivers in falls prevention cannot be overlooked. Education and support for families are essential components of effective falls prevention strategies. Geriatric nurses can provide guidance to caregivers on how to create a safe living environment and encourage their loved ones to engage in preventive measures. This collaborative approach not only enhances the safety of older adults but also fosters a supportive network that can contribute to their overall well-being [7].

As we explore the various evidence-based interventions available for falls prevention and mobility support, it is crucial to emphasize the importance of individualized care. Each older adult is unique, with distinct health profiles, living situations, and personal preferences. Therefore, interventions must be tailored to meet the specific needs of each individual. This personalized approach not only increases the likelihood of successful outcomes but also empowers older adults to take an active role in their health and safety [8].

### **Understanding Falls in the Elderly**

To effectively prevent falls, it is essential to understand the multifaceted nature of the problem. Falls among older adults can result from a combination of intrinsic factors, which are internal to the individual, and extrinsic factors, which are external environmental influences. Intrinsic factors include age-related physiological changes that affect an individual's ability to maintain balance and coordination. These changes encompass a range of physical and cognitive impairments, such as muscle weakness, balance impairments, cognitive decline, and visual deficits. As people age, their muscle mass and strength naturally decline, which can lead to decreased mobility and an increased risk of falling. The loss of muscle strength, particularly in the lower extremities, can significantly impact an individual's stability and ability to perform daily activities [9].

Cognitive decline, which may manifest as dementia or other forms of cognitive impairment, further

complicates the risk of falls. Individuals with cognitive impairments may struggle with spatial awareness, judgment, and the ability to assess risks in their environment, making them more susceptible to falls. Visual deficits, such as cataracts or macular degeneration, can hinder an older adult's ability to see obstacles or changes in elevation, leading to an increased likelihood of tripping or losing balance. Additionally, chronic conditions such as arthritis, diabetes, and cardiovascular diseases can further exacerbate these risks. Arthritis can limit mobility and range of motion, while diabetes may lead to neuropathy, affecting balance and coordination. Cardiovascular diseases can contribute to dizziness and fainting spells, particularly when transitioning from sitting to standing, creating a perfect storm for falls [10].

Extrinsic factors encompass environmental hazards that can contribute to falls. Poor lighting, slippery floors, uneven surfaces, and cluttered living spaces are common environmental factors that increase the likelihood of falls. For instance, inadequate lighting in hallways or stairwells can make it difficult for older adults to navigate safely. Similarly, slippery floors due to spills or wax can create hazardous conditions, particularly for those with compromised balance. Uneven surfaces, such as cracked sidewalks or thresholds, pose additional risks, as they can easily catch a foot and lead to a fall. Cluttered living spaces can also obstruct pathways, increasing the chances of tripping over objects that are not easily visible. Understanding the interplay between intrinsic and extrinsic factors is vital for geriatric nurses as they assess the risk of falls in their patients and develop tailored interventions. By recognizing how these factors interact, nurses can create comprehensive strategies that address both the individual's physical condition and their environment, ultimately reducing the risk of falls [11].

### **Risk Assessment**

Effective falls prevention begins with a comprehensive risk assessment. Geriatric nurses are uniquely positioned to conduct thorough assessments that identify both intrinsic and extrinsic risk factors. Utilizing standardized assessment tools, such as the Morse Fall Scale or the Hendrich II Fall Risk Model, nurses can systematically evaluate a patient's risk of falling based on their medical history, physical condition, and environmental factors. These tools provide a structured approach to risk assessment, allowing for the identification of high-risk individuals who may benefit from targeted interventions [12].

In addition to standardized assessments, nurses should engage in direct observation of patients' mobility and functional abilities. This observational approach allows nurses to identify specific challenges that patients may face in their daily activities, such as difficulty navigating stairs or getting in and out of bed. By watching how patients move and interact with their environment, nurses can gain valuable insights into their functional capabilities and limitations [13]. For example, a patient who struggles to maintain balance while walking may require a different intervention than one who has difficulty with spatial awareness. By combining quantitative assessments with qualitative observations, nurses can create a comprehensive picture of a patient's fall risk and develop targeted interventions that address the specific needs of each individual [14].

Furthermore, the assessment process should not be a one-time event. Continuous monitoring and re-evaluation of fall risk are essential, particularly as patients age or as their health status changes. Geriatric nurses should establish a routine for re-assessing fall risk at regular intervals or whenever there is a significant change in a patient's health condition. This ongoing evaluation allows for timely adjustments to care plans and interventions, ensuring that they remain effective and relevant to the patient's current situation [15].

### **Evidence-Based Interventions**

Numerous evidence-based interventions have been identified to reduce the incidence of falls among older adults. These interventions can be categorized into several key areas, including exercise programs, environmental modifications, medication management, and education. Each of these areas plays a crucial role in creating a comprehensive falls prevention strategy that addresses the diverse needs of older adults [16].

### **Exercise Programs**

Regular physical activity is one of the most effective strategies for preventing falls in older adults. Exercise programs that focus on improving strength, balance, and flexibility have been shown to significantly reduce fall risk. Research indicates that engaging in balance training exercises can lead to a substantial decrease in falls among older adults. Tai Chi, for instance, is a low-impact exercise that emphasizes balance and coordination, making it particularly beneficial for older individuals. Studies have demonstrated that participation in Tai Chi can lead to a 30% reduction in fall risk

among older adults, making it a highly recommended activity for this population. The slow, controlled movements of Tai Chi not only enhance physical stability but also promote mindfulness, which can further reduce anxiety related to falling [17].

Resistance training is another effective intervention that can enhance lower body strength and stability. Geriatric nurses can facilitate access to community-based exercise programs or develop individualized exercise plans for patients. These plans should be tailored to the individual's current physical capabilities and health conditions, ensuring that they are both safe and effective. Encouraging regular physical activity not only improves physical function but also promotes mental well-being, reducing the fear of falling and enhancing overall quality of life. Additionally, group exercise programs can foster social interaction, which is beneficial for mental health and can motivate older adults to remain active [18]. Moreover, incorporating flexibility exercises into physical activity regimens can further enhance mobility and reduce the risk of falls. Stretching exercises can improve range of motion and help maintain joint health, which is particularly important for older adults who may experience stiffness or discomfort. Geriatric nurses should emphasize the importance of a well-rounded exercise program that includes strength, balance, and flexibility training to maximize the benefits for their patients [19].

### **Environmental Modifications**

Environmental modifications are essential components of falls prevention strategies. Simple changes in the home environment can significantly reduce fall risk. Nurses can conduct home safety assessments to identify potential hazards and recommend modifications. These may include removing clutter, securing loose rugs, improving lighting, and installing grab bars in bathrooms. Each of these modifications addresses specific risks that can lead to falls, creating a safer living environment for older adults [20]. In addition to home modifications, community environments should also be assessed for safety. Geriatric nurses can advocate for safer public spaces by collaborating with local authorities to ensure that sidewalks are well-maintained, public restrooms are accessible, and community centers offer appropriate activities for older adults. For instance, ensuring that public spaces are well-lit and free of obstacles can encourage older adults to engage in outdoor activities, which can further enhance their physical health and social well-being [21]. By addressing environmental factors, nurses can create safer spaces for older adults to live and engage in social activities, ultimately contributing to their overall quality of life. Furthermore, educating patients and their families about the importance of maintaining a safe home environment is crucial. Nurses can provide resources and guidance on how to make necessary modifications, empowering families to take an active role in falls prevention. This collaborative approach not only enhances safety but also fosters a sense of community and support among caregivers and older adults [22].

### **Medication Management**

Polypharmacy, or the use of multiple medications, is common among older adults and is a significant risk factor for falls. Certain medications, particularly sedatives, antihypertensives, and antidepressants, can contribute to dizziness, confusion, and impaired coordination. Geriatric nurses play a vital role in medication management by conducting thorough medication reviews and collaborating with healthcare providers to optimize pharmacotherapy. This process involves evaluating the necessity of each medication, considering potential side effects, and identifying any drug interactions that may increase fall risk [23].

Regular medication reviews can help identify potentially inappropriate medications and facilitate dose adjustments or discontinuation where necessary. Additionally, educating patients about the side effects of their medications and the importance of adherence can enhance safety and reduce fall risk. Nurses can also encourage patients to communicate openly with their healthcare providers about any concerns related to their medications. This proactive approach to medication management not only reduces the risk of falls but also promotes overall health and well-being among older adults [24].

Moreover, geriatric nurses should be aware of the importance of non-pharmacological interventions for managing conditions that may otherwise require medication. For example, lifestyle modifications, such as dietary changes and increased physical activity, can often improve health outcomes without the need for additional medications. By advocating for these alternatives, nurses can help reduce the overall medication burden on older adults, further decreasing the risk of falls [25].

### **Education and Empowerment**

Education is a powerful tool in falls prevention. Geriatric nurses can provide education to older adults and their caregivers about the importance of falls prevention strategies. This education can encompass topics such as the significance of regular exercise, the need for home safety modifications, and the role of medication management in preventing falls.

Empowering older adults to take an active role in their health can lead to improved outcomes.

Nurses can encourage patients to engage in discussions about their health, set personal goals for mobility, and participate in decision-making regarding their care. By fostering a sense of autonomy and responsibility, nurses can help older adults feel more confident in their ability to prevent falls [26].

### **Multidisciplinary Collaboration**

Falls prevention requires a collaborative approach involving multiple healthcare professionals. Geriatric nurses can work alongside physical therapists, occupational therapists, pharmacists, and physicians to develop comprehensive care plans tailored to individual needs. This multidisciplinary collaboration ensures that all aspects of falls risk are addressed, from physical function to medication management [27].

Regular communication among team members is essential for effective intervention. Interdisciplinary meetings can facilitate the sharing of insights and strategies, allowing for a more holistic approach to patient care. By leveraging the expertise of various professionals, geriatric nurses can enhance the effectiveness of falls prevention strategies and improve patient outcomes [28].

### **Conclusion**

Falls prevention and mobility support are critical components of geriatric nursing that require a multifaceted approach. By understanding the risk factors associated with falls, conducting thorough assessments, and implementing evidence-based interventions, geriatric nurses can significantly reduce the incidence of falls among older adults. The integration of exercise programs, environmental modifications, medication management, and education empowers older adults to take control of their health and enhances their quality of life. As the population continues to age, the role of geriatric nursing in falls prevention will become increasingly important. By fostering collaboration among healthcare professionals and advocating for safe environments, geriatric nurses can lead the charge in creating a safer, more supportive community for older adults. Ultimately, the goal is to promote independence, enhance mobility, and reduce the risk of falls, ensuring that older adults can live fulfilling and active lives.

---

### **References:**

1. (2021). Effects of the nursing intervention fall prevention in older adults with arterial hypertension using nanda-i, nic, and noc. *international journal of nursing knowledge*, 33(2), 147-161. <https://doi.org/10.1111/2047-3095.12346>
2. (2019). Caring for the older person with cognitive impairment in hospital: qualitative analysis of nursing personnel reflections on fall events. *journal of clinical nursing*, 28(7-8), 1346-1353. <https://doi.org/10.1111/jocn.14724>
3. (2011). Incidence and cost of serious fall-related injuries in nursing homes. *clinical nursing research*, 21(1), 10-23. <https://doi.org/10.1177/1054773811414180>
4. (2020). Bedrails and falls in nursing homes: a systematic review. *clinical nursing research*, 30(1), 5-11. <https://doi.org/10.1177/1054773820907805>
5. (2012). Patient safety and falls: a qualitative study of home care nurses in norway. *Nursing and health sciences*, 14(4), 452-457. <https://doi.org/10.1111/j.14422018.2012.00701.x>
6. (2017). Nurses' perceptions of preventing falls for patients with dementia in the acute hospital setting. *australasian journal on ageing*, 36(4). <https://doi.org/10.1111/ajag.12474>
7. (2020). Home and environmental hazards modification for fall prevention among the elderly. *public health nursing*, 38(3), 493-501. <https://doi.org/10.1111/phn.12852>
8. (2012). Barriers to senior centre implementation of falls prevention programmes. *injury prevention*, 18(4), 272-276. <https://doi.org/10.1136/injuryprev-2011-040204>
9. (2021). Nurses' perceptions of recommended fall prevention strategies. *journal of nursing care quality*, 37(3), 249-256. <https://doi.org/10.1097/ncq.0000000000000605>
10. (2023). Risk of falls and strategies to reduce fall rates in nursing homes.. <https://doi.org/10.21203/rs.3.rs-2989885/v1>
11. (2011). Randomized controlled trial comparing tailoring methods of multimedia-based fall prevention education for community-dwelling older adults. *american journal of occupational therapy*, 65(6), 702-709. <https://doi.org/10.5014/ajot.2011.001180>
12. (2018). Management team perceptions of risks and strategies for preventing falls among short-stay patients in nursing homes. *the health care manager*, 37(1), 76-85.

<https://doi.org/10.1097/hcm.00000000000000192>

13. (2015). Factors associated with increased hospitalisation risk among nursing home residents in sweden: a prospective study with a three-year follow-up. *international journal of older people nursing*, 11(2), 130-139. <https://doi.org/10.1111/1468-2317.12107>
14. (2017). Falls prevention process in assisted living communities. *journal of applied gerontology*, 38(6), 805-824. <https://doi.org/10.1177/0733464817748776>
15. (2013). Risk factors for falls in older people in nursing homes and hospitals. a systematic review and meta-analysis. *archives of gerontology and geriatrics*, 56(3), 407-415. <https://doi.org/10.1016/j.archger.2012.12.006>
16. (2016). Impact of fall prevention on nurses and care of fall risk patients. *the gerontologist*, gnw156. <https://doi.org/10.1093/geront/gnw156>
17. (2012). Falls in newly admitted nursing home residents: a national study. *journal of the American geriatrics society*, 60(5), 939-945. <https://doi.org/10.1111/j.15325415.2012.03931.x>
18. (2017). Us and dutch nurse experiences with fall prevention technology within nursing home environment and workflow: a qualitative study. *geriatric nursing*, 38(4), 276-282. <https://doi.org/10.1016/j.gerinurse.2016.11.005>
19. (2011). Environmental assessment and modification to prevent falls in older people. *Journal of the American geriatrics society*, 59(1), 26-33. <https://doi.org/10.1111/j.15325415.2010.03221.x>
20. (2017). Barriers and enablers to the implementation of the 6-pack falls prevention program: a pre-implementation study in hospitals participating in a cluster randomised controlled trial. *plos one*, 12(2), e0171932. <https://doi.org/10.1371/journal.pone.0171932>
21. (2020). Nursing intervention assessment tool fall prevention in elderly people with systemic arterial hypertension. *journal of nursing education and practice*, 10(7), 60. <https://doi.org/10.5430/jnep.v10n7p60>
22. (2017). Opinions of nurses about the evaluation of risk of falling among inpatients. *rehabilitation nursing*, 42(6), E19-E24. <https://doi.org/10.1002/rnj.281>
23. (2011). Reduction of femoral fractures in long-term care facilities: the bavarian fracture prevention study. *plosone*, 6(8), e24311. <https://doi.org/10.1371/journal.pone.0024311>
24. (2019). Evaluation of respond, a patient-centred program to prevent falls in older people presenting to the emergency department with a fall: a randomised controlled trial. *plos medicine*, 16(5), e1002807. <https://doi.org/10.1371/journal.pmed.1002807>
25. (2014). Shared medical appointments to screen for geriatric syndromes: preliminary data from a quality improvement initiative. *journal of the american geriatrics society*, 62(12), 2415-2419. <https://doi.org/10.1111/jgs.13142>
26. (2018). Views of key stakeholders on the causes of patient falls and prevention interventions: a qualitative study using the international classification of functioning, disability and health. *journal of clinical nursing*, 28(3-4), 15-628. <https://doi.org/10.1111/jocn.14656>
27. (2020). Characteristics and circumstances of falls in the community-dwelling older adult population. *journal of primary care & community health*, 11. <https://doi.org/10.1177/2150132720940508>
28. (2021). Effectiveness of educational interventions for fall prevention: a systematic review. *texto & contexto - enfermagem*, 30. <https://doi.org/10.1590/1980-265x-tce2020-0558>